

Heavy Axions from Strong Broken Horizontal Gauge Symmetry

T. Elliott and S. F. King

Physics Department

University of Southampton

Southampton

SO9 5NH

U.K.

Abstract

We study the consequences of the existence and breaking of a Peccei–Quinn symmetry within the context of a dynamical model of electroweak symmetry breaking based on broken gauged flavour symmetries. We perform an estimate of the axion mass by including flavour instanton effects and show that, for low cut-offs, the axion is sufficiently massive to prevent it from being phenomenologically unacceptable. We conclude with an examination of the strong CP problem and show that our axion cannot solve the problem, though we indicate ways in which the model can be extended so that the strong CP problem is solved.

1 Introduction

The origin of electroweak symmetry breaking remains an unsolved problem. While the Higgs sector of the standard model may well be an accurate reflection of the underlying physical processes driving electroweak symmetry breaking, and simultaneously giving quarks and leptons their masses, it is possible that it is only an approximation to some underlying dynamical mechanism.

The realisation that the top quark, if it exists, must have a large mass has led many authors to speculate about the role of the top quark in electroweak symmetry breaking [1]. In particular, it is possible that a top quark condensate, driven by some as yet undiscovered dynamics, is responsible for electroweak symmetry breaking. In order to produce a condensate one needs to induce a 4-quark contact term with a sufficiently large coupling:

$$\mathcal{L} = \mathcal{L}_{kin} + G_t(\bar{Q}_L t_R)(\bar{t}_R Q_L), \quad (1)$$

where $Q_L^T = (t_L, b_L)$. Recently several authors have begun to consider the dynamics which may allow such an interaction to appear in the low energy effective theory [2].

If we wish to allow the bottom quark to form a condensate, as surely we must if we wish to maintain maximal correspondence between u,c,t quarks and d,s,b quarks (and, besides, it will transpire that, in the models we consider, we cannot give mass to the d,s,b sector any other way), it is necessary to introduce the additional interaction

$$G_b(\bar{Q}_L b_R)(\bar{b}_R Q_L) \quad (2)$$

However, since we shall admit this term, we may include all gauge-invariant interactions possible including only the top and bottom quarks [3].

$$\begin{aligned} \mathcal{L} = \mathcal{L}_{kin} &+ G_t(\bar{Q}_L t_R)(\bar{t}_R Q_L) + G_b(\bar{Q}_L b_R)(\bar{b}_R Q_L) \\ &+ G_{tb} \left((\bar{Q}_L b_R)(\bar{t}_R^c \tilde{Q}_L) + h.c. \right), \end{aligned} \quad (3)$$

where $\tilde{Q} = i\sigma_2 Q^c$, and c denotes charge conjugation. If the contact operators arise from heavy gauge boson exchange [2], then $G_{tb} \equiv 0$, since the only gauge boson which can mediate $b_L \rightarrow t_L$ transitions is the W . This leads to a problem as discussed below.

It is easy to show that the Lagrangian density (3) is entirely equivalent to the following, in which the 4-fermi couplings are reparametrised as Higgs fields without kinetic terms (the kinetic terms are radiatively induced)

$$\begin{aligned}\mathcal{L} = \mathcal{L}_{kin} \quad & - \quad \mu_t^2 \phi_t^\dagger \phi_t + (\bar{Q}_L \tilde{\phi}_t t_R + h.c.) \\ & - \quad \mu_b^2 \phi_b^\dagger \phi_b + (\bar{Q}_L \phi_b b_R + h.c.) \\ & - \quad \mu_{tb}^2 (\phi_t^\dagger \phi_b + h.c.)\end{aligned}\tag{4}$$

If $\mu_{tb} \equiv 0$, then the model is the standard two Higgs doublet model with a $U(1)$ Peccei–Quinn (PQ) symmetry [4]

$$\begin{aligned}Q_L &\rightarrow e^{i\alpha} Q_L & t_R &\rightarrow e^{-i\alpha} t_R & b_R &\rightarrow e^{-i\alpha} b_R \\ \phi_t &\rightarrow e^{-2i\alpha} \phi_t & \phi_b &\rightarrow e^{2i\alpha} \phi_b\end{aligned}\tag{5}$$

Thus, upon spontaneous symmetry breakdown, not only does $SU(2)_L \otimes U(1)_Y$ get broken to $U(1)_{em}$, but also the ungauged PQ symmetry breaks resulting in a Goldstone boson. However, the PQ symmetry is explicitly broken by QCD instanton effects [5], and the pseudoscalar becomes the standard electroweak axion [4, 6]. A light pseudoscalar with mass ~ 20 keV is ruled out experimentally. This is a serious problem for dynamical theories in which the contact operators are generated by heavy gauge boson exchange.

In what follows we examine a recently proposed model based on broken gauged chiral flavour symmetries [7]. It will be seen that this model also possesses multiple Higgs doublets and a PQ symmetry. We show, however, that the PQ symmetry is broken not only by QCD instantons, but also by instantons associated with the gauged flavour groups and some technicolour-like groups. (Henceforth we shall refer to the technicolour-like groups as ultracolour groups). Since these groups will be strongly coupling at some scale Λ , with $\Lambda \geq 1$ TeV, it will be seen that, under certain circumstances, the axion will acquire a mass sufficiently large in order to prevent it from being phenomenologically unacceptable.¹

In section 2 we describe the models we shall consider. In section 3 we calculate estimates of the axion mass in these models and discuss the implications of its cut-off

¹Without reference to a particular model, Yamawaki [8] conjectured that instanton effects arising from gauged flavour symmetries might provide another source of explicit PQ symmetry breaking, therefore raising the axion mass from its standard QCD value.

dependence. Finally, in section 4, we describe the possible role of this axion in solving the strong CP problem, and conclude that it does not. We describe extensions of the models in which the strong CP problem is solved.

2 “Moose” Models

One set of dynamics, which has been proposed recently, is based on a gauged flavour symmetry which is broken by non-quark condensates driven by ultracolour interactions [7]. These condensates in turn allow the formation of quark condensates resulting in both dynamically generated quark masses and electroweak symmetry breaking.

In the models which we will consider a multiplicity of both gauged and ungauged symmetry groups will proliferate. It is therefore convenient, at this stage, to describe a notation, of which we shall make extensive use, in which the particle content and particle transformation properties are completely transparent [9]. An $SU(n)$ group is represented by the number n . If the number is circled, then the group is gauged; otherwise it is a global symmetry. Directed lines correspond to left-handed Weyl fermions. A line attached to a number, either circled or uncircled, indicates that the fermions transform under that group. If the line is directed away, then the fermions transform under the fundamental representation; if directed towards the number, then the fermions transform under the conjugate representation.

For simplicity we restrict ourselves at first to an idealised world in which the down, strange and bottom quarks are absent. We then postulate the existence of an ultracolour interaction which acts on new left-handed Weyl fermions, and the gauging of flavour. The new symmetry groups, where ‘ga’ means gauged and ‘gl’ means global, are

$$SU(3)_c \otimes SU(3)_f^{ga} \otimes SU(3)_L^{ga} \otimes SU(3)_R^{ga} \otimes SU(3)_L^{gl} \otimes SU(3)_R^{gl} \quad (6)$$

and the quarks $U_L^T \equiv (u_L, c_L, t_L)$, $U_R^{cT} \equiv (u_R^c, c_R^c, t_R^c)$ and new fermions a, b, c, d

transform as

$$\begin{aligned}
U_L &\sim (3 , 1 , 3 , 1 , 1 , 1) \\
U_R^c &\sim (\bar{3} , 1 , 1 , \bar{3} , 1 , 1) \\
a &\sim (1 , 3 , \bar{3} , 1 , 1 , 1) \\
b &\sim (1 , \bar{3} , 1 , 3 , 1 , 1) \\
c &\sim (1 , 3 , 1 , 1 , 1 , \bar{3}) \\
d &\sim (1 , \bar{3} , 1 , 1 , 3 , 1)
\end{aligned} \tag{7}$$

$SU(3)_{L,R}^{ga}$ are gauged chiral flavour symmetries acting on the left- and right-handed quarks, respectively. $SU(3)_c$ is the QCD gauge group. We may introduce a mass matrix connecting the c and d particles. Providing that the elements of the matrix are small compared to the typical dynamical masses, the explicit breaking of the global symmetries associated with the c and d particles will be slight. In figure 1 we give the “moose” corresponding to this model. It is obvious that such a notation enables us to see immediately the structure of the theory.

We now assume that $SU(3)_f^{ga}$ becomes strongly coupling at some scale Λ_f and drives the formation of the following condensates

$$\begin{aligned}
&< a_i d_i > \neq 0 , < b_i c_i > \neq 0 , i = 1, 2 \\
&< a_3 b_3 > \neq 0 , < c_3 d_3 > \neq 0
\end{aligned} \tag{8}$$

Here we do not discuss the precise nature of the vacuum alignment calculations, but simply refer the interested reader to a recent analysis [10]. It is sufficient to state that if we wish to form a top quark condensate, then we need a large value of $\tilde{M}_{33}^U$, the mass matrix element connecting the c_3 and d_3 particles, though not sufficiently large to break badly the global chiral symmetry. Given such a value, however, calculations reveal that the condensates of equation (8) are preferred over

$$< a_i d_i > \neq 0 , < b_i c_i > \neq 0 , i = 1, 2, 3 \tag{9}$$

The preferred condensates of (8) break the groups $SU(3)_L^{ga} \otimes SU(3)_R^{ga}$ and $SU(3)_L^{gl} \otimes SU(3)_R^{gl}$ down to the diagonal subgroup $SU(2)_L \otimes SU(2)_R$ — the global symmetry of the standard model in the case of a very massive top quark. Of course, the symmetry will be broken by the quark mass matrix. That we are able to break the model down to the standard model together with its associated global symmetries ensures that we have a G.I.M. mechanism [11, 9], the absence of which, hitherto, has been the bane of technicolour theories. A G.I.M. mechanism is essential if Λ_f is to be lowered below about 10^3 TeV, so that flavour-changing neutral-currents are suppressed.

The Goldstone bosons generated by the breaking of the gauged and global groups become the longitudinal components of the gauge bosons associated with the $SU(3)_L^{ga}$ and $SU(3)_R^{ga}$ groups, thereby rendering them massive. In the limit $g_{L,R} \rightarrow 0$ (the coupling constants associated with the groups $SU(3)_L^{ga}$ and $SU(3)_R^{ga}$, respectively) we may calculate the gauge boson mass matrix, here 16 by 16, which, in general, requires diagonalisation to determine the mass eigenstates [7]. Mixing between the L- and R-sectors can be so introduced. It is not necessary here to give the form of the gauge boson mass matrix. Indeed, it is more edifying, as we shall see, to work with the gauge eigenstates.

Associated with, for example, the a and d particles there is also a $U(1) \otimes U(1)$ symmetry, an axial symmetry. The condensates will break this down to the diagonal subgroup, $U(1)$, generating yet another Goldstone boson. However the $U(1)$ symmetries are anomalous [12] and thus the (pseudo-)Goldstone boson receives a mass because of $SU(3)_f$ instanton effects [5]. The dynamics of this mechanism are identical to those whereby the η acquires a mass $O(\Lambda_{QCD})$ because of the explicit breaking of the $U(1)_A$ quark symmetry by QCD instantons. Thus, our analogue, the “ultra- η ” will have a mass $O(\Lambda_f)$.

The condensates described above thus allow us to induce interactions between the left- and right-handed quarks. Working with the gauge boson gauge eigenstates, the lowest order contribution to a t_L - t_R interaction comes from the diagram of figure 2. A cross denotes a condensate insertion. Of course, there are similar diagrams for c_L - c_R and u_L - u_R interactions except that the $(T^8)_{33}$ is replaced by $(T^8)_{11}$ or $(T^8)_{22}$, T^a being the generators of $SU(3)$. Thus, such diagrams give contributions 4 times smaller than the t_L - t_R interaction.

Such a diagram induces an interaction which, after Fierz re-arrangement, is of the form

$$g_L g_R A (\bar{t}_L t_R) (\bar{t}_R t_L), \quad (10)$$

where A is some number with dimensions $[M]^{-2}$ determined by group theory and the dynamics of the inner loop [7]. This is precisely the term we require in order now to drive the formation of a top quark condensate, breaking electroweak symmetry and giving the top quark a large dynamical mass. We must adjust g_L and g_R so that we

obtain critical chiral symmetry breaking. That being so, because the corresponding 4-fermion interactions of up and charm quarks have an overall coefficient 4 times smaller than that for the top quark, we will not obtain up and charm quark condensates in addition.

In fact, to form a top quark condensate the flavour groups must be strongly coupling, though broken, in addition to the ultracolour groups. Thus, it is natural to be concerned about the possibility that the strongly coupling flavour groups will themselves drive other, unwanted, condensates, for example $\langle a_i U_i \rangle \neq 0$, which would break the ultracolour groups and QCD. We simply assume that this does not occur.

Masses for the up and charm quarks are derived from the top mass by diagrams such as those in figure 3. A blob represents the insertion of a mass matrix element $(\tilde{M}^U)_{ab}$ connecting the c and d particles. The corresponding interactions, again after Fierz rearrangements, are

$$\begin{aligned} & (\tilde{M}^U)_{22} B(\bar{t}_L t_R)(\bar{c}_R c_L) \\ & (\tilde{M}^U)_{11} B(\bar{t}_L t_R)(\bar{u}_R u_L) \end{aligned} \quad (11)$$

If $\langle \bar{t}t \rangle \neq 0$, we see that the up and charm quarks acquire a mass proportional to $(\tilde{M}^U)_{ii} \langle \bar{t}t \rangle$.

To extend these ideas to include the down, strange and bottom quarks is straightforward. Of course, the left-handed quarks must be placed into $SU(2)_L$ doublets. Thus, we essentially obtain a moose which corresponds to 2 copies of the moose of figure 1, except that the flavour groups acting on the left-handed quarks are compressed into one flavour group acting on the electroweak doublets, Q_L

$$Q_L^T = \left(\begin{pmatrix} u \\ d \end{pmatrix}_L, \begin{pmatrix} c \\ s \end{pmatrix}_L, \begin{pmatrix} t \\ b \end{pmatrix}_L \right). \quad (12)$$

Notice that the right-handed quarks U_R^c transform under a different flavour group from the right-handed quarks $D_R^{cT} = (d_R^c, s_R^c, b_R^c)$. The moose is in figure 4. (That which has been described above applies equally to the right-hand side of the moose of figure 4.)

The moose of figure 4, however, has a non-trivial vacuum alignment calculation associated with it. We do not discuss the problem, but, again, refer the reader to a

recent analysis [10]. The result of the analysis, however, is that the mass matrices connecting both the c and d and the g and h particles, from which the quark mass matrices descend, can be simultaneously diagonalised. Thus, we obtain no understanding of the structure of the KM mass matrix. We must therefore conclude that the mass matrices themselves descend from some new dynamics not included in the present moose.

Calculations performed elsewhere reveal that one has to take Λ_f very large in order to obtain a top quark mass compatible with constraints on the ρ -parameter [7]. Naively, of course, if a particle obtains a dynamical mass from dynamics at the scale Λ , then the particle should acquire a mass $O(\Lambda)$. To prevent such an occurrence one has to fine-tune. We believe, as a general principle, that fine-tuning should be avoided if possible. Thus, we should like to take $\Lambda_f \sim 1$ TeV, this being the lowest scale possible because of phenomenological constraints. If a quark condensate is responsible for electroweak symmetry breaking it is not unreasonable to consider the possibility of a fourth generation of quarks and leptons. We may then take $\Lambda_f \sim 1$ TeV, since there is as yet no phenomenological limit on the masses of degenerate, fourth generation quarks. Such low cut-offs, moreover, would enable the models to be falsifiable at, for example, the S.S.C. We shall call the fourth generation quarks T and B.

The relevant fourth generation moose is readily obtainable from that of figure 4, simply by replacing the $SU(3)$ flavour groups by $SU(4)$ flavour groups, and the $SU(3)$ global groups by $SU(4)$ global groups.

3 Calculation of the Axion Mass

We have seen that interactions of the form $(\bar{Q}_L t_R)(\bar{t}_R Q_L)$ and $(\bar{Q}_L b_R)(\bar{b}_R Q_L)$ can be obtained from the moose model. Reflection upon the structure of the moose in figure 4 reveals that we can never induce the term

$$G_{tb} \left((\bar{Q}_L b_R)(\bar{t}_R^c \tilde{Q}_L) + h.c. \right) \quad (13)$$

in our low energy theory. The two operators $(\bar{t}_L b_R)(\bar{t}_R^c b_L^c)$ and $(\bar{b}_L b_R)(\bar{t}_R^c t_L^c)$ would descend from the un-Fierzed $(\bar{t}_L \gamma^\mu b_L^c)(\bar{t}_R^c \gamma_\mu b_R)$ and $(\bar{b}_L \gamma^\mu t_L^c)(\bar{t}_R^c \gamma_\mu b_R)$, and these clearly

can never arise from diagrams of the type in figure 4. Thus $G_{tb} \equiv 0$ and the PQ symmetry is explicitly broken by instantons only.

Each line (or particle) in the moose of figure 4 transforms under not only the groups to which the line is attached, but also a $U(1)$ group. We have already indicated that the non-quark condensates break $U(1)$ symmetries, but since these symmetries are anomalous, we obtain “ultra- η ” particles with large mass. Associated with the three different quark lines we have the three $U(1)$ symmetries $U(1)_{D_R} \otimes U(1)_Q \otimes U(1)_{U_R}$; these symmetries can be rewritten in the more familiar form of $U(1)_B \otimes U(1)_Y \otimes U(1)_{PQ}$. $U(1)_B$ is unbroken by the quark condensates; $SU(2)_L \otimes U(1)_Y$ is broken down to $U(1)_{em}$, with a neutral Goldstone boson which becomes the longitudinal component of Z^0 . $U(1)_{PQ}$ is ungauged, and its breaking results in a (pseudo-)Goldstone boson, the axion [4, 6].

We limber up for the calculation of the axion mass in the moose model by reminding the reader of how to calculate its mass in QCD. This will enable us to develop further familiarity with the moose notation and hence allow us to see immediately with which kind of processes flavour or ultracolour instanton amplitudes are associated. We consider two flavour QCD, $N_f = 2$. The moose appropriate for this case is that shown in figure 5. The effective QCD instanton vertex is shown in figure 6 and is associated with a process in which $\Delta Q_5 = 2N_f$, that is, the axial charge changes by $2N_f$ units, and the vacuum tunnels to a state of different Pontryagin index [5].

In order to obtain a non-zero contribution to the vacuum energy density, one must close off the instanton zero modes. In this way, by coupling the fermions produced in a tunnelling process to some external, chirality flipping particles, such as Higgs scalars, one may obtain the diagram of figure 7. Of course, this diagram explicitly breaks the PQ symmetry, rendering the axion massive. The amplitude for this process is given by [5]

$$\begin{aligned} \mathcal{M} &= g_u g_d K e^{-8\pi^2/g^2(\Lambda_{QCD})} \int \frac{d\rho}{\rho^5} \rho^{3N_f} (\rho \Lambda_{QCD})^\beta \\ &\times \prod_{i=1,2} \int d^4x \frac{2}{\pi^2} \phi_i(x) \frac{1}{[\rho^2 + x^2]^3}, \end{aligned} \quad (14)$$

where g_u and g_d are the couplings of ϕ_1 to u and ϕ_2 to d, respectively, and K is given

by

$$K = \frac{4}{\pi^2} \left(\frac{4\pi^2}{g^2(\Lambda_{QCD})} \right)^6 \frac{1}{2} e^{A-B}, \quad (15)$$

where $A = 7.054$, $B = 0.360$ and $\beta = 11 - \frac{2}{3}N_f$. It is to be understood that there is present a factor $e^{i\theta}$, θ being the QCD θ -parameter. Evaluating the integrals (we cut the spatial integrals off at the lower end with ρ , and the instantons with sizes $\sim 1/\Lambda_{QCD}$ are assumed to dominate), and using Dashen's formula [13] to give us the mass of a pseudo-Goldstone boson

$$(f_a m_a)^2 = \langle 0 | [Q^a, [Q^a, \mathcal{M}]] | 0 \rangle, \quad (16)$$

where Q^a is the charge associated with the axial current, and f_a is the axion decay constant ($f_a \approx 250$ GeV if the PQ symmetry is spontaneously broken at the same scale as electroweak symmetry breaking), we obtain

$$(f_a m_a)^2 \sim m_u m_d \Lambda_{QCD}^2 \quad (17)$$

Of course, this is precisely the result we expect from dimensional analysis and the η mass.

Comparison of the structure of the QCD moose in figure 5 with the moose of figure 4, together with knowledge of the effective vertex to which QCD instantons give rise, allows us to see by inspection the nature of the processes to which the flavour and ultracolor instantons give rise. It is easy to see that the flavour and ultracolor groups together provide another source of PQ symmetry breaking. $SU(3)_U$ instantons force U_R^c and a to rotate together axially; $SU(3)_{fU}$ instantons force a and b to rotate together, and so on. Thus, from the point of view of explicit PQ symmetry breaking, the whole moose may be regarded as equivalent to one in which the quarks transform under two non-abelian groups — instantons from both sectors break the PQ symmetry.

Of course, the flavour groups are dynamically broken, and the gauge bosons become massive. Finite action, non-zero solutions of the Euclidean equations of motion cease to exist since classical scale invariance is broken, and we can always rescale non-zero solutions and reduce the action. However, in Euclidean space, approximate solutions, near which the action varies slowly, still dominate. The effect of the breaking of classical scale invariance is to suppress the constrained instanton [14] solutions

with a Yukawa decay at distances larger than the inverse mass scale. We therefore encode this behaviour by cutting off the integrals over instanton size at the upper end by $1/\Lambda$, Λ being the scale at which the symmetry is broken. However, if the gauge groups are strongly coupling at the scale Λ , then we may still obtain significant contributions to instanton-induced amplitudes.

The Higgs particles in our dynamical model are, of course, composite objects, bound states of quarks. Since left- and right-handed quarks transform under different flavour groups, and as there are no current insertions which can change t_L into b_{3R} , there is no one-instanton contribution to the axion mass. The minimal diagram contains three different instantons, where, in effect, we have closed off the zero modes of an $SU(3)_Q$ instanton on $SU(3)_U$ and $SU(3)_D$ instantons. However, we now keep the discussion general and suppose that we have n generations of quarks and leptons. The diagram is shown in figure 8. The large hatched blobs, of course, represent the effective vertex of an instanton amplitude; the small unhatched circles denote a composite Higgs coupling to a fermion; r, g, b distinguish colours, with tilded colours labelling $SU(3)_{fU}$ colours and doubly tilded colours labelling $SU(3)_{fD}$ colours. The e_3 and b_3 particles are understood to be right-handed, and the a_3 and f_3 are understood to be left-handed. The cross, as usual, denotes a condensate insertion. Inserting condensates is acceptable since we may suppose that chiral symmetry breaking occurs before confinement, as is supposed for QCD.

In estimating the contribution of this rather complicated process we assume that we may perform a naive perturbation theory in the effective instanton vertices. This naive assumption seems to be vindicated by recent analyses of the restoration of unitarity in scattering processes by multi-instanton contributions to the total scattering amplitude [15] and, moreover, by the seminal work of Callen, Dashen and Gross [5]. Provided that we work in the dilute gas approximation, any ambiguity regarding the zero modes of the total vacuum tunnelling amplitude is removed, and each sub-process has a clearly defined number of zero modes.

To calculate the amplitude we need to perform the integral

$$I = \frac{2}{\pi^2} \int d^4y \frac{1}{[\rho_x^2 + (y-x)^2]^{\frac{3}{2}}} \frac{1}{[\rho_z^2 + (y-z)^2]^{\frac{3}{2}}}, \quad (18)$$

where we have replaced the Higgs fields by their vacuum expectation values in an-

anticipation of employing Dashen's formula. We obtain an angular integral which is a complete elliptic integral of the second kind. Since we require only an order of magnitude estimate of the axion mass, we approximate the integrand by its value at $\theta = \pi/2$. We thus obtain

$$I \approx \frac{4}{[s^2 + \rho_z^2 - \rho_x^2]^2} \left(\sqrt{s^2 + \rho_z^2} - \rho_x \right)^2, \quad (19)$$

where $s^2 = (z - x)^2$. Recalling that we are working in the dilute gas approximation, and that the ρ integrals are cut off, $s \gg \max\{\rho_x, \rho_z\}$, we may simply take

$$I \approx \frac{4}{s^2} \quad (20)$$

and integrate s from $1/\Lambda_U$ to ∞ , Λ_U being the scale at which $SU(n)_U$ is broken. Since the groups $SU(3)_U$, $SU(3)_D$ and $SU(3)_Q$ are all broken at approximately the same scale, we set $\Lambda_U = \Lambda_D = \Lambda_Q = \Lambda$. We assume, moreover, that the gauge couplings g_U , g_D and g_Q are approximately equal (to g). There are six such legs emerging from the $SU(n)_U$ instanton and six from the $SU(n)_D$ instanton, so we need,

$$J = \left[\int_{1/\Lambda} d^4s \left(\frac{4}{s^2} \right)^6 \right]^2 = 2^{20} \pi^4 \Lambda^{16} \quad (21)$$

For each instanton we must perform an integration over its size, with a cut-off corresponding to the scale at which instantons become suppressed because the group is broken. We need

$$\sigma_D = \sigma_U = \int_0^{1/\Lambda} \frac{d\rho}{\rho^5} \rho^{3.3} (\rho\Lambda)^{(\frac{11}{3}n - \frac{2}{3}3)}, \quad (22)$$

and

$$\sigma_Q = \int_0^{1/\Lambda} \frac{d\rho}{\rho^5} \rho^{3.6} (\rho\Lambda)^{(\frac{11}{3}n - \frac{2}{3}6)}. \quad (23)$$

So,

$$\sigma_U = \sigma_D = \frac{3}{11n + 9} \Lambda^{-5} \quad (24)$$

$$\sigma_Q = \frac{3}{11n + 30} \Lambda^{-14} \quad (25)$$

This completes all the integrations, so we may put the result together:

$$\begin{aligned} (f_a m_a)^2 &= K_U K_D K_Q \left(e^{(-8\pi^2/g^2)} \right)^3 \\ &\times (m_U m_D)^3 (\Lambda_{fU} \Lambda_{fD})^3 \\ &\times \frac{27}{(11n + 9)^2 (11n + 30)} \Lambda^{-8} 2^{20} \pi^4, \end{aligned} \quad (26)$$

where $D=b$ or B ($n=3$ or 4), $U=t$ or T ($n=3$ or 4), and

$$K_i = \frac{4}{\pi^2} \left(\frac{4\pi^2}{g^2} \right)^{2n} \frac{1}{(n-1)!(n-2)!} e^{A+B(n-2-N_f^i)}, \quad (27)$$

$i = U, D$ or Q , and $N_f^U = N_f^D = 3$, $N_f^Q = 6$. Λ_{fU} and Λ_{fD} are taken as the values of the condensate insertions. We suppose that the scales at which the ultracolour groups break chiral symmetry are approximately equal, and that $\Lambda_{fU} \approx \Lambda_{fD} = \Lambda$. Extracting the n dependent terms from this result, together with the dimensionful quantities, and calling the remaining constant C , we have

$$(f_a m_a)^2 = C \frac{e^{3nB}}{(11n+9)^2(11n+30)} \frac{1}{[(n-1)!(n-2)!]^3} \frac{m_D^3 m_U^3}{\Lambda^2} \quad (28)$$

It is well-known that instanton calculations are rather unreliable. Thus we seek to determine C from general physical principles. The axion receives a mass from dynamics at the scales associated with the flavour and ultracolour groups, but arises as a (pseudo-)Goldstone boson because of spontaneous symmetry breaking at the weak scale. Now, if the ultracolour groups become strongly coupling at ~ 1 TeV, so that the associated ultra- Π decay constants are ~ 250 GeV — the electroweak scale — we expect to have a theory with no fine-tuning and thus one in which all particle masses are ~ 1 TeV. Of course, for the top quark, such a mass is unacceptably large. However, for the T and B quarks, 1 TeV is perfectly reasonable. The axion will obtain a mass from dynamics at the scale at which the PQ symmetry is spontaneously broken. Thus, provided that the couplings of the Higgs particles to the quarks are of order unity, as they must be since $m_T = g_T < \phi_T >$ and $m_T \approx < \phi_T > \approx 1$ TeV, the axion would also have a mass ~ 1 TeV. (The axion mass in the standard model is so small because $f_\pi \ll f_a$, and the up and down quarks are very light.) We may use this general argument to determine C . Taking $f_a = 250$ GeV, all mass scales ~ 1 TeV, and $n = 4$ (4 generations) we obtain

$$C = 3.0 \times 10^5 \quad (29)$$

This value of C corresponds to taking $\alpha_Q = \alpha_U = \alpha_D \approx 7$ at a scale of 1 TeV. This agrees with our expectation that the flavour groups need to be strongly coupling in order to drive quark condensates. Calculations performed elsewhere [9] indicate that $\alpha \approx 30$ is required to obtain quark condensates. However, these calculations are

based on a truncation of a Dyson–Schwinger equation and thus should be regarded as indicating the general qualitative features of the theory, rather than providing reliable estimates of physically interesting quantities.

For $n = 3$ and $n = 4$, we have

$$f_a m_a = \frac{2.9}{\Lambda} \sqrt{m_t^3 m_b^3} \quad n = 3 \quad (30)$$

$$f_a m_a = \frac{1}{4\Lambda} \sqrt{m_T^3 m_B^3} \quad n = 4 \quad (31)$$

Taking $m_t = 125$ GeV and $m_b = 5$ GeV, for $\Lambda = 1$ TeV, we obtain $m_a = 183$ MeV for 3 generations of quarks, and for $m_T = m_B = 1$ TeV, $m_a = 1$ TeV for 4 generations of quarks, as arranged.

Clearly $m_a = 1$ TeV is acceptable, and possibly $m_a = 183$ MeV is also acceptable phenomenologically [16]. In the above estimates we have assumed $\Lambda = 1$ TeV, and for the three generation model we have taken $m_t = 125$ GeV rather than the much larger value expected from standard calculations [1]. If these standard calculations are at all reliable, then we would expect to have a much larger value of Λ in the three generation case, for a reasonable value of m_t . For example, $m_t \approx 230$ GeV requires $\Lambda \approx 10^{15}$ GeV [1]. For such a large value of Λ it is clear from equation (30) that the axion receives a contribution to its mass from flavour instantons of $f_a m_a \sim 10^{-10} (\text{GeV})^2$, which is much too small. Thus, our calculations rule out the possibility of three families ² with a top quark condensate induced by heavy gauge boson exchange, due to an unacceptably light axion.

4 The Strong CP Problem

The standard model possesses two possible sources of strong CP violation: the QCD θ -parameter; phases in the quark mass matrix. In the absence of colour instantons one may re-define the quark fields so that the mass matrix is real, and in the absence of a quark mass matrix one may re-define the quark fields and set $\theta = 0$. However, in the presence of instantons and phases in the quark mass matrix, there is a potential

²Interestingly there is another problem with the three family model due to the decay $\Upsilon \rightarrow \gamma H'_0$, where H'_0 is a Higgs scalar associated with the $\langle \bar{b}b \rangle$ condensates. In the three generation model with $G_{tb} = 0$ we expect $m_{H'_0} < 2m_b$, which is apparently ruled out by the above process.

conflict between the direction chosen for zero CP violation on the vacuum manifold. This conflict would result in overall strong CP violation. Experimentally it is known that $\theta + \arg \det M = \bar{\theta} \leq 10^{-9}$. This is the strong CP problem. For a review of the strong CP problem see, for example, [16] and references therein.

In the two Higgs doublet extended standard model, the Peccei–Quinn–Wilczek–Weinberg axion provides a natural understanding of the absence of strong CP violation. Through its coupling to the anomaly as $\frac{a}{32\pi^2 f_a} F_{\mu\nu}^a \tilde{F}_{\mu\nu}^a$, and after re-defining the axion field by $a \rightarrow a - f_a \bar{\theta}$, one can show that the vacuum of smallest energy density corresponds to $\langle a \rangle = 0$. Thus, the axion forces the two directions of zero CP violation on the vacuum manifold to be consistent. Unfortunately there is no empirical evidence to suggest that the PQWW axion exists.

Given the occurrence of multiple Higgs doublets in the present model, and a massive axion, it is natural to ask whether the strong CP problem is solved here. We begin by demonstrating that there is only one overall θ -parameter in the moose of figure 4, rather than 6.

We may work at a sufficiently high energy scale (temperature) above which the vacuum has undergone a phase transition and entered a phase in which all the condensates have evaporated. In this phase, all particles are massless, except that the c–d and g–h mass matrices remain because they are non-dynamical. Because all dynamical masses vanish, we are at liberty to perform axial re-definitions of the particle fields. Consider, for example, the U_R^c line which has associated with it two θ -parameters, θ_c , the colour θ -parameter, and θ_U , the flavour θ -parameter of the right-handed u,c,t (t) quarks. Now we perform an axial re-definition of the U_R^c fields. Because the functional measure is not invariant under such a re-definition [17], both θ_c and θ_U change. If we change the phase of the U_R^c fields by α , then $\theta_c \rightarrow \theta_c - 2n\alpha$ and $\theta_U \rightarrow \theta_U + 2n_c\alpha$ (n is the number of generations, n_c the number of quark colours). Setting $2n\alpha = \theta_c$ we may rotate θ_c to zero at the expense of changing θ_U to $\theta_U + \frac{n_c}{n}\theta_c$. By rotating other fields we may move all θ -parameters onto, for example, the $SU(3)_{fU}$ group. All other θ -parameters would be zero and thus would give no CP violation. The θ -parameter associated with the $SU(3)_{fU}$ group would be $\theta'_{fU} = \theta_{fU} + \theta_{fD} + \theta_c + \frac{n_c}{n}(\theta_Q + \theta_U + \theta_D)$, in an obvious notation. Moreover, by performing axial re-definitions of the c and d fields, we may rotate θ'_{fU} to zero and place it, instead, as a phase in the c–d mass

matrix. Any other configurations of θ -parameters, experimentally, must be identical to that which we have just obtained.

Our axion is a composite of quarks, the quarks feeling only colour and flavour forces. Therefore the axion does not couple to ultracolour fields. In particular, the axion cannot be used to argue that the state of minimum vacuum energy density occurs when $\bar{\theta}_{fU} = \theta'_{fU} + \arg \det M_{cd} = 0$. Thus, there will always exist CP violating effects associated with the ultracolour groups. Moreover, because the c–d mass matrix contains CP violating phases, and since the quark mass matrices are proportional to the c–d and g–h mass matrices, strong CP violation will be re-introduced when the vacuum enters the phase in which condensates form. Consequently, the present model provides no solution to the strong CP problem.

There are, however, three variants of the present model which would enable us to solve the strong CP problem, the last being extremely speculative:

Variant 1:

We have no real understanding of the c–d and g–h mass matrices — we have to assume that they descend from some other set of dynamics. We could simply impose that the ultrastrong dynamics respects CP symmetry. Thus, because the ultrastrong dynamics respect CP, so would QCD.

Variant 2:

We now have precisely the QCD situation over again: c and d transform under $SU(3)_{fU}$, we have a mass matrix and ultrastrong CP violation. From the absence of strong CP violation we know that there can be no ultrastrong CP violation. Any dynamics giving c and d their mass may be reparametrised, for convenience, by a Higgs sector. In particular, therefore, we may have another PQ symmetry. Thus c and d acquire their masses at some scale ≥ 1 TeV, and the new PQ symmetry relaxes ultrastrong CP violation to zero. A new axion would exist, but would be massive because $\Lambda_{fU} \geq 1$ TeV (as opposed to $\Lambda_{QCD} \sim 250$ MeV), and, moreover, would not couple to quarks and leptons. We would not expect to be aware of its existence until we probed the dynamics of the c–d mass generation mechanism.

Variant 3:

There is also an alternative solution which relies on the fact that neutrinos are very

light, if not massless. We know that, if the up quark were massless, strong CP violation would not exist, since all CP violating effects could be rotated onto the up quark mass term which, of course, would be zero. If we could find a way of dumping all CP violation onto the neutrino mass term, we would similarly have a solution. To include leptons within the current formalism, however, presents some considerable difficulty [9, 18]. The construction and viability of the quark moose critically relies on the fact that quarks come in more than one colour. Gauge anomaly cancellation in the flavour groups then requires us to have an $SU(n_c)$ group for the ultracolour groups. This is a wonderful coincidence, for we require that the ultracolour groups are non-abelian in order that they may be asymptotically free. Unfortunately each flavour of lepton comes in only one ‘colour.’ Thus, we need strongly coupling $U(1)$ groups [9] !

In order to circumvent this difficulty, and in order to establish quark–lepton symmetry, we suppose that leptons come in three colours [19, 7]. The $SU(3)$ of colour on the leptons is broken at some high scale to $SU(2)$. We then allow the $SU(2)$ colour group to become confining, so that the leptons transforming under the $SU(2)$ subgroup of the original $SU(3)$ of colour on the leptons do not appear in the low energy spectrum of particles. The remaining leptons, those transforming under the broken generators of the $SU(3)$ colour group, are precisely those which one sees at low energy. Henceforth we shall simply refer to unbroken lepton colour, but with the understanding that there is a mechanism for breaking lepton colour. The lepton moose, then, is entirely equivalent to the quark moose of figure 4. Notice that quark–lepton symmetry demands the existence of right-handed neutrinos. A fourth generation neutrino would have to be massive so as not to conflict with recent LEP data. Since the u, c, t (and T) quark masses are parametrised as the mass matrix connecting the c and d particles, let the neutrino masses be parametrised as a mass matrix connecting the c' and d' particles, where the c' and d' particles are the lepton moose analogue of the quark moose c and d particles.

If we could establish a link between the quark and lepton moose, we could then solve the strong CP problem. The simplest manner in which to establish a connection is to demand that the ultracolour groups belonging to the quark moose and the lepton moose be identical. Anomaly cancellation tells us that the number of lepton colours

equals the number of quark colours. The $c'-d'$ mass matrix is tiny, if not identically zero, since the neutrinos are very light. In the 4 generation model, only $(M_{c'd'})_{44}$ would be non-zero, so as to allow the fourth generation neutrino to become massive. We may use our previous analysis to argue that all θ -parameters can be moved onto $SU(3)_{fU}$, together with any phases in any of the mass matrices. Now we exploit the massless neutrino by rotating the overall $\bar{\theta}$ onto the (non-existent or small) $c'-d'$ mass terms. Hence CP violating effects can be sucked away, and the massless neutrino solves the CP problem. This conclusion is relatively model independent, needing only a connection between the source of quark and lepton masses via non-abelian groups.

5 Conclusions

Any dynamical theory of electroweak symmetry breaking and mass generation based on gauged, chiral horizontal symmetries possesses a PQ symmetry which is spontaneously broken at the weak scale. The appearance and breaking of the PQ symmetry is, at first sight, a disaster, for one would assume that the resulting pseudoscalar is the standard electroweak axion, which is ruled out experimentally. We have seen, however, that the explicit breaking of the PQ symmetry by flavour instantons provides a contribution to the axion mass over and above the usual contribution from colour instantons. The calculations performed above reveal that the axion mass is inversely proportional to the scale characteristic of the new flavour and ultracolour dynamics. Therefore, for large Λ , the flavour instanton contribution to the axion mass is greatly suppressed, as decoupling would demand. *This means that any dynamical theory based on principles similar to those described above, and which requires a large Λ , is ruled out, due to the non-existence of the standard axion.* However, all theories to date suggest that Λ must be large if the top quark mass is to be consistent with constraints on the ρ -parameter. The simplest way to avoid this conflict is to assume the existence of a fourth generation of quarks and leptons. Λ may then be ~ 1 TeV, the T and B quarks give no contribution to the ρ -parameter as they are mass degenerate, and, moreover, the axion becomes sufficiently heavy to prevent it from being a problem phenomenologically. These conclusions are relatively model independent. Particular to our model, however, is the presence of a G.I.M. mechanism which al-

lows us to lower Λ below $\sim 10^3$ TeV without fear of phenomenological difficulties associated with flavour-changing neutral-currents.

Unfortunately, the present model does not solve the strong CP problem: in order to understand the absence of strong CP violation, we need to understand the absence of ultrastrong CP violation. We have indicated some ways in which such an understanding could be obtained. The most speculative extension imposes a quark-lepton symmetry, and the masslessness of the neutrinos supplies us with a natural understanding of the absence of ultrastrong and therefore strong CP violation.

Acknowledgements

S.F.K. is grateful to the S.E.R.C. for the support of an Advanced Fellowship, and T.E. is grateful to the S.E.R.C. for the support of a studentship.

References

- [1] Y Nambu, “New Theories in Physics,” in Proc XI Warsaw Symposium on Elementary Particle Physics (ed Z Ajduk *et al*, World Scientific, Singapore, 1989).
V A Miransky, M Tanabashi, K Yamawaki, Phys Lett **B221**, (1989), 177.
W A Bardeen, C T Hill, M Lindner, Phys Rev **D41**, (1990), 1647.
W J Marciano, Phys Rev Lett **62**, (1989), 2793.
- [2] S F King, S H Mannan, Phys Lett **B241**, (1990), 249.
M Lindner, D A Ross, Nucl Phys **B370**, (1992), 30.
C T Hill, Phys Lett **B266**, (1991), 419.
- [3] M A Luty, Phys Rev **D41**, (1990), 2893.
M Suzuki, Phys Rev **D41**, (1990), 3457.
- [4] R Peccei, H Quinn, Phys Rev Lett **38**, (1977), 1440 ; Phys Rev **D16**, (1977), 1791.
- [5] A A Belavin, A M Polyakov, A S Schwartz, Yu S Tyupkin, Phys Lett **59B**, (1975), 85.
G ’t Hooft, Phys Rev Lett **37**, (1976), 8 ; Phys Rev **D14**, (1976), 3432.
C G Callen, R G Dashen, D J Gross, Phys Lett **63B**, (1976), 334.
R Jackiw, C Rebbi, Phys Rev Lett **37**, (1976), 172.
- [6] S Weinberg, Phys Rev Lett **40**, (1978), 223.
F Wilczek, Phys Rev Lett **40**, (1978), 279.
- [7] S F King, Phys Rev **D45**, (1992), 990.
T Elliott, S F King, “Heavy Quark Condensates from Dynamically Broken Flavour Symmetry,” Southampton preprint SHEP 91/92–13 (1992), to appear in Phys Lett B.
- [8] K Yamawaki, “Top–Mode Standard Model,” in Proc 1990 International Workshop on Strong Coupling Gauge Theories and Beyond (ed T Muta, K Yamawaki, World Scientific, Singapore, 1991).

- [9] H Georgi, “Technicolour and Families,” in Proc 1990 International Workshop on Strong Coupling Gauge Theories and Beyond (ed T Muta, K Yamawaki, World Scientific, Singapore, 1991).
- [10] N J Evans, S F King, D A Ross, “Top Quark Condensation from Broken Family Symmetry,” Southampton preprint SHEP 91/92–11 (1992).
- [11] S L Glashow, J Iliopoulos, L Maiani, Phys Rev **D2**, (1970), 1285.
- [12] J S Bell, R Jackiw, Nuovo Cimento **60A**, (1969), 47.
S L Adler, Phys Rev **117**, (1969), 2426.
- [13] R Dashen, Phys Rev **183**, (1969), 1245.
- [14] I Affleck, Nucl Phys **B191**, (1981), 429.
- [15] H Aoyama, H Kikuchi, Phys Rev **D43**, (1991), 1999.
- [16] H–Y Cheng, Phys Reports **158**, (1988), 1.
- [17] K Fujikawa, Phys Rev Lett **42**, (1979), 1195 ; Phys Rev **D21**, (1980), 2848.
- [18] R S Chivukula, H Georgi, H L Randall, Nucl Phys **B292**, (1987), 93.
R S Chivukula, H Georgi, Phys Lett **B188**, (1987), 99.
S. Dimopoulos, H Georgi, S Rabi, Phys Lett **127B**, (1983), 101.
- [19] R Foot, H Lew, Phys Rev **D41**, (1990), 3502.

Figure Captions

Figure 1 : The toy moose. The $SU(3)_c$ group is, of course, colour.

Figure 2 : A ‘bubble’ graph causing interactions between t_L and t_R^c , ultimately giving rise to a top quark condensate. The cross represents a condensate insertion.

Figure 3 : ‘Bubble’ graphs allowing up and charm quarks to acquire a mass from the top quark mass. The blob denotes a mass insertion.

Figure 4 : Two moose bolted together, for four families.

Figure 5 : The simple QCD moose.

Figure 6 : A QCD effective instanton vertex explicitly breaking the $U(1)_A$ quark symmetry.

Figure 7 : An QCD instanton–induced, PQ violating contribution to the axion potential.

Figure 8 : The minimal instanton–induced, PQ violating contribution to the axion potential in the moose model. x , y , and z are the centres of the instantons, and ρ_x , ρ_y , and ρ_z are the instanton sizes.